

Review History, Presenting Symptoms, and Risk Factors. Identify as Low/Intermediate/High Risk per Guidelines for TOLAC

Low Risk

- * CNM in House
- * OB in House

- * No More than 1-2 previous Cesareans
- * Spontaneous Labor
- * Category I Tracing

Intermediate Risk

- * CNM on L&D
- * OB readily available
- * OB aware of risk level
- * OB informed of changes in status and reassesses patient
- * Anesthesia has interviewed patient and documented interview
- * Anesthesia available in-house
- * Non-prostaglandin method of Induction/Augmentation acceptable
- * Any Induction/Augmentation of Labor
- * Category II Tracing

High Risk

- * CNM on L&D – no other patient care responsibilities
- * OB on L&D – no other patient care responsibilities
- * Anesthesia available to L&D – no other patient care responsibilities
- * No change in dilation/descent after two hours of adequate labor
- * Bleeding not considered “cervical show”
- * Category III Tracing

ALL TOLACS:

- Provider review of consent** with customer owner; Confirmed desire for TOLAC; Provider note documented upon admission
- Anesthesia** notification of TOLAC/VBAC and assessment documented on admission
- All patients undergoing TOLAC have:**
 - Large Bore IV in place (Consider placing 2nd Large Bore IV)
 - Type and Screen; CBC
 - Continuous Monitoring in active labor
 - Option for Epidural
 - Provider with privileges for Cesarean delivery In-House