

Oral Titrated Misoprostol for Induction of Labor at Alaska Native Medical Center (ANMC)

Table of Contents

BACKGROUND	2
METHODS FOR INDUCTION OF LABOR	2
MISOPROSTOL DOSING	2
MISOPROSTOL CONTRAINDICATIONS AND CAUTIONS	3
CERVICAL RIPENING/INDUCTION WITH LOW DOSE TITRATED ORAL MISOPROSTOL AT ANMC	3
CANDIDATES	3
CONTRAINDICATIONS	3
ORAL MISOPROSTOL INDUCTION OF LABOR PROCEDURE	4
ORAL MISOPROSTOL LABOR AUGMENTATION PROCEDURE	5
UTERINE TACHYSYSTOLE DURING MISOPROSTOL ADMINISTRATION	5
TRANSITIONING FROM MISOPROSTOL TO OXYTOCIN	5
REFERENCES	6
VERSION HISTORY	9
TABLES	10
TABLE 1: BISHOP SCORE	10
TABLE 2: CERVICAL RIPENING – ORAL TITRATED MISOPROSTOL PROTOCOL	10
TABLE 3: AUGMENTATION – ORAL TITRATED MISOPROSTOL PROTOCOL	11
FIGURES	12
FIGURE 1: LEVEL OF PLASMA MISOPROSTOL CONCENTRATION OVER TIME (ZEIMAN, ET AL).	12

BACKGROUND

The incidence of labor induction has been steadily rising over the past decades. The rate of induced labor reached 34.5% in the United States during 2024, which is an increase of 32-67% from 2016. Specifically in Alaska, the percent change in induction of labor rate from 2016 (26.1%) to 2024 (36.8%) was 41%. There are numerous fetal or maternal medical indications for delivery. Induction of labor (IOL) without a medical indication may occur on/after 39 0/7 weeks gestation if the patient has been thoroughly counseled about the risks and benefits of labor induction versus expectant management.

The increased rate of labor induction is likely multifactorial in nature, but a large study, the ARRIVE Trial (A Randomized Trial of Induction Versus Expectant Management), also likely contributed to this increase. The ARRIVE Trial was a large, randomized controlled trial over 41 institutions with 94% adherence that evaluated the risks and benefits of IOL in low-risk nulliparous women at 39 weeks of gestation. The findings demonstrated decreased rates of cesarean delivery, decreased rates of hypertensive disorders, no difference in composite adverse neonatal outcomes, and better outcomes for both mothers and babies. The American College of Obstetricians and Gynecologists (ACOG) supports using these findings to guide shared decision making and accommodate elective inductions of labor.

METHODS FOR INDUCTION OF LABOR

Induction of labor can be a long process with considerable resource expenditure. The Bishop score (Table 1) is a clinical tool to determine if a cervix is favorable or “ripe” for induction. If the score is five or less, the cervix is unfavorable and cervical ripening agents should be used. If the score is eight or higher, the cervix is favorable and there is a high chance of a vaginal delivery.

Cervical ripening increases the likelihood of a vaginal birth within 24 hours when compared with use of oxytocin alone. There are mechanical and pharmacologic agents that are effective at cervical ripening. There are many different mechanical options such as a Foley balloon, Atad balloon, Cook cervical ripening balloon, seaweed-based dilators (Laminaria), and osmotic agents (Dilapan). Pharmacologic methods primarily include prostaglandins such as misoprostol (Cytotec) and dinoprostone (Cervidil), although occasionally oxytocin (Pitocin) is used in select candidates. Review of the current evidence suggests that the prostaglandin misoprostol, administered either orally or intravaginally, is generally considered the most effective agent for cervical ripening. Misoprostol has a good safety profile at the lower dosage range, and is convenient to use, but there are limitations to its use. Although safe and used commonly for labor induction, misoprostol use for cervical ripening is currently off label.

MISOPROSTOL DOSING

The optimal dose of misoprostol for labor induction is 25 mcg or less, administered orally or vaginally. As misoprostol is supplied in 100 or 200 mcg tablets, dosing in less than 50 mcg fragments may be inaccurate and inconsistent. Likewise, misoprostol may not be evenly distributed within the inert material of the tablet, making failure of response, or an exaggerated response, unpredictable.

In response to this challenge, dissolving a 200 mcg tablet in 200 mL of water is proposed as an accurate, safe, and efficacious method of using the drug in lower doses. The dosing interval of every two hours for

oral misoprostol is based on the known pharmacokinetics of the drug. When misoprostol is administered orally, it requires 30 to 60 minutes to achieve peak plasma levels, but is then promptly metabolized, and is essentially undetectable after 120 minutes. (Figure 1)

MISOPROSTOL CONTRAINDICATIONS AND CAUTIONS

Misoprostol use is contraindicated in those that have a hypersensitivity reaction to its components, history of uterine rupture, or history of prior full thickness myometrial surgery such as cesarean delivery or myomectomy. Misoprostol can be associated with uterine tachysystole, which may result in fetal heart rate abnormalities, an effect which is both dose and route of administration dependent.

CERVICAL RIPENING/INDUCTION WITH LOW DOSE TITRATED ORAL MISOPROSTOL AT ANMC

CANDIDATES

1. Women with a cephalic singleton pregnancy, no contraindications to misoprostol, and with an unfavorable cervix (Bishop score ≤ 5) who are:
 - a. Preterm ($\leq 36\ 6/7$ weeks) with a medical indication for delivery or
 - b. Full term ($\geq 37\ 0/7 - 39\ 0/7$ weeks) with a medical indication for delivery or
 - c. Full term ($\geq 39\ 0/7$ weeks) with or without a medical indication for delivery (elective).
2. Women with a breech singleton pregnancy, no contraindications to misoprostol, an unfavorable cervix (Bishop score ≤ 5), an OB provider skilled in vaginal breech delivery, and appropriate counseling with shared decision making.
3. Women with a multifetal gestation (dichorionic/diamniotic twin pregnancy or monochorionic/diamniotic twin pregnancy), cephalic presenting fetus, no contraindications to misoprostol, have a provider skilled in multifetal deliveries who previously counseled the patient and is available for the induction and delivery process, and with an unfavorable cervix (Bishop score ≤ 5).

CONTRAINDICATIONS

1. Women who are NOT candidates for this protocol include those with:
 - a. Prior full-thickness myometrium uterine surgery (cesarean delivery or myomectomy).
 - b. Non-reassuring antenatal fetal surveillance (i.e. significant recurrent decelerations of the fetal heart rate noted during fetal monitoring).
 - c. Active labor: more than three palpable contractions lasting greater than 30 seconds in 10 minutes, or cervical dilation greater than 4 cm.
 - d. Any known obstetric contraindication to labor (i.e. placenta previa, vasa previa, breech fetal position without an OB provider skilled in vaginal breech delivery, active genital herpes simplex infection, etc).
 - e. Hypersensitivity reaction to misoprostol or its components.

ORAL MISOPROSTOL INDUCTION OF LABOR PROCEDURE

1. In the pharmacy, misoprostol 200 mcg is dissolved in 200 mL of tap water, constituting a solution of 20 mcg in 20 mL. This solution may be used for up to 48 hours. (USP 795)
2. Induction of labor medication regimen:
 - a. 20mcg given orally every one-hour x 4 doses.
 - b. 40mcg given orally every two hours x 4 doses.
 - c. 60mcg given orally every two hours x 4 doses.
3. Once the patient is admitted to Labor & Delivery and orders have been placed, obtain vital signs, perform a sterile vaginal exam, confirm fetal presentation, and perform a 20-minute fetal non-stress test (NST).
 - a. If any of the following occurs, the patient is NOT considered a ripening candidate with oral misoprostol:
 - i. Nonreactive NST with or without decelerations
 - ii. Bishop score ≥ 6
 - iii. Regular painful uterine contractions every 3 minutes lasting 30 seconds
 - iv. Active labor
 - v. Contraindications to a vaginal delivery (see above)
4. An initial dose of **misoprostol 20 mcg** (20 mL measured in a syringe) will be given by mouth, and the time recorded by the nurse.
 - a. The same dose will be repeated every hour for three hours (total of four doses).
 - b. Patients may ambulate, take a bath, eat, and drink during the ripening procedure.
 - c. The patient will be monitored with an NST and tocography 20 minutes after receiving the medication. If fetal status remains reassuring and there are no other indications for monitoring, then the patient can be off the monitor for 20 minutes. The patient is then placed back on the monitor 20 minutes prior to the next dose.
5. If the FHR remains reassuring, and regular uterine contractions (3 contractions every 10 minutes lasting at least 30 seconds) have *not* ensued after 4 hours:
 - a. The misoprostol dose may be increased to **40 mcg** (40 mL) every two hours until active labor occurs.
 - b. Again, the patient will be monitored with an NST and tocography for 20 minutes after receiving the medication. If fetal status remains reassuring and there are no other indications for monitoring, then the patient can be off the monitor until 20 minutes prior to the next dose.
 - c. If not previously performed, a sterile vaginal exam should be performed 12 hours after the first dose of misoprostol has been given.
6. If active labor has still not begun in 12 hours, the misoprostol dose may be increased to **60 mcg** (60 mL) every two hours as long as the fetal heart rate pattern remains reassuring and other contraindications have not occurred.
 - a. Similarly, the patient will be monitored with an NST and tocography for 20 minutes after receiving the medication. If fetal status remains reassuring and there are no other indications for monitoring, then the patient can be off the monitor until 20 minutes prior to the next dose.
 - b. If not previously performed, a sterile vaginal exam should be performed 24 hours after the first dose of misoprostol. If active labor has not yet begun, the provider will decide on the patient's management.

ORAL MISOPROSTOL LABOR AUGMENTATION PROCEDURE

1. If active labor is established, but contractions become inadequate (less than 3 contractions in 10 minutes lasting 30 seconds over a 2-hour period), or labor is arrested (no further dilation over 2 hours), and the patient declines other augmentation methods, then the labor may be **augmented** with titrated oral misoprostol
 - a. Beginning at 5 mcg every hour, progressing to 10 mcg every hour, or 20 mcg every hour (maximum dose permitted) every hour until adequate labor progress is established.
 - b. The patient will be monitored with cardiotocography for 20 minutes after receiving the medication. If fetal tolerance has been assured, then the patient can be off the monitor for 20 minutes. The pt is then placed back on the fetal heart tracing and cardiotocogram (CTG) 20 minutes prior to the next dose.

UTERINE TACHYSYSTOLE DURING MISOPROSTOL ADMINISTRATION

1. Definition: more than 5 contractions in 10 minutes, averaged over a 30-minute window.
2. Uterine tachysystole **without** worrisome fetal heart rate changes, is not an indication to stop the misoprostol procedure.
3. Uterine tachysystole **with** fetal heart rate abnormalities (Category II or III tracings) should result in holding the next misoprostol dose and discussing the plan of care with the provider. Routine resuscitative measures such as maternal repositioning or fluid boluses can be given per unit guidelines.

TRANSITIONING FROM MISOPROSTOL TO OXYTOCIN

1. Timeline:
 - a. Wait at least one hour after administering 20mcg misoprostol orally.
 - b. Wait at least two hours after administering either 40mcg or 60mcg orally.

REFERENCES

1. Martin JA, Osterman JK. Induction of Labor Increases in the United States: 2016 – 2024. National Center for Health Statistics Data Brief #554. March 25, 2026.
2. Management of Full-Term Nulliparous Individuals Without a Medical Indication for Delivery: ACOG Clinical Practice Update. *Obstetrics & Gynecology* 145(1):p e45-e50, January 2025.
3. ACOG. Induction of labor. ACOG Practice Bulletin No. 107. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2009;114: 386–97. (Reaffirmed 2020)
4. Aalami-Harandi R, Karamali M, Moeini A. Induction of labor with titrated oral misoprostol solution versus oxytocin in term pregnancy: randomized controlled trial. *Rev Bras Ginecol Obstet*. 2013 Feb;35(2):60-5.
5. Abramovici D, Goldwasser S, Mabie BC, Mercer BM, Goldwasser R, Sibai BM. A randomized control trial of oral misoprostol versus Foley catheter and oxytocin for induction of labor at term. *Am J Obstet Gynecol* 1999; 181:1108-12.
6. Alfirevic Z, Keeney E, Dowswell T, Welton NJ, Medley N, Dias S, Jones LV, Caldwell DM. Methods to induce labour: a systematic review, network meta-analysis and cost-effectiveness analysis. *BJOG*. 2016 Aug;123(9):1462-70
7. Alfirevic Z, Keeney E, Dowswell T, Welton NJ, Dias S, Jones LV, Navaratnam K, Caldwell DM. Labour induction with prostaglandins: a systematic review and network meta-analysis. *BMJ*. 2015 Feb 5;350:h217.
8. Hung CH, Cheng SY, Chan TM, Lee MR. Evaluating misoprostol content in pregnant women with hourly oral administration during labor induction by microElution solid phase extraction combined with liquid chromatography tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2015 Sep 1;1000:176-80
9. Berard V, Fiala C, Cameron S, Bombas T, Parachini M, Gemzell-Danielsson K. Instability of misoprostol tablets stored outside the blister: a potential serious concern for clinical outcome in medical abortion. *PLoS One*. 2014 Dec 15;9(12):e112401.
10. Bricker L, Peden H, Tomlinson AJ, Al-Hussaini TK, Idama T, Candelier C, Luckas M, Furniss H, Davies A, Kumar B. Titrated low-dose vaginal and/or oral misoprostol to induce labour for prelabour membrane rupture: a randomised trial. *BJOG*. 2008;115:1503-1511.
11. Cheng SY, Hsue CS, Hwang GH, Chen W, Li TC. Comparison of labor induction with titrated oral misoprostol solution between nulliparous and multiparous women. *J Obstet Gynaecol Res*. 2010 Feb;36(1):72-8.
12. Cheng SY. How to Manage Labor Induction or Augmentation to Decrease the Cesarean Deliveries Rate, Cesarean Delivery, Dr. Raed Salim (Ed.), 2012. ISBN: 978-953-51-0638-8, (Accessed 4/15/23)
13. Cheng SY. Individualized misoprostol dosing for labor induction or augmentation: A review. *World J Obstet Gynecol* 2013; 2(4): 80-86 <http://www.wjnet.com/2218-6220/full/v2/i4/80.htm#B23> (Accessed 4/15/23).
14. Cheng SY, Ming H, Lee JC. Titrated oral compared with vaginal misoprostol for labor induction: a randomized controlled trial. *Obstet Gynecol*. 2008;111:119-125.
15. Cheng SY, Chen TC. Pilot study of labor induction with titrated oral misoprostol. *Taiwan J Obstet Gynecol*. 2006;45:225-229
16. Dallenbach P, Boulvain M, Viardot C, Irion O. Oral misoprostol or vaginal dinoprostone for labor induction: a randomized controlled trial. *Am J Obstet Gynecol* 2003;188:162-7.

17. Dodd JM, Crowther CA, Robinson JS Oral misoprostol for induction of labour at term: randomised controlled trial. *BMJ* 2006;332;509-513;
18. Elati A, Week AD. The use of misoprostol in obstetrics and gynecology. *BJOG* 2009; 116(S1):61-9.
19. French L. Oral prostaglandin E2 for induction of labour. *Cochrane Database Syst Rev* 2001; CD00309. (Accessed 4/15/23)
20. Ho M, Cheng SY, Li TC. Titrated oral misoprostol solution compared with intravenous oxytocin for labor augmentation: a randomized controlled trial. *Obstet Gynecol.* 2010 Sep;116(3):612-8.
21. Hofmeyr GJ, Alfirevic Z, Matonhodze B, et al. Titrated oral misoprostol solution for induction of labor: a multicenter randomized trial. *BJOG* 2001; 108:952-9.
22. Hofmeyr GJ, Matonhodze BB, Alfirevic Z, Campbell E, de Jager M, Nikodem C. Titrated oral misoprostol solution--a new method of labour induction. *S Afr Med J.* 2001 Sep;91(9):775-6.
23. How HY, Leaseburg L, Khoury JC, Siddi I TA, Spinnato JA, Sibai BM. A comparison of various routes and dosages of misoprostol for cervical ripening and induction of labor. *Am J Obstet Gynecol* 2001; 185:911-5.
24. Kundodyiwa TW, Alfirevic Z, Weeka AD. Low dose oral misoprostol for induction of labor: a systematic review. *Obstet Gynecol* 2009; 113:374-83.
25. Majoko F, Zwizwai M, Nystrom L, Lindmark G (2002) Vaginal misoprostol for induction of labour: a more effective agent than prostaglandin f2 alpha gel and prostaglandin e2 pessary. *Central African Journal of Medicine.* pp. 123-128.
26. Matonhodze BB, Hofmeyr GJ, Levin J. Labour induction at term: a randomized control trial comparing Foley catheter and titrated oral misoprostol solution, titrated oral misoprostol solution alone, and dinoprostone. *South African Med J* 2003; 93:375-9.
27. Moodley J, Venkatachalam S, Songca P. Misoprostol for cervical ripening at and near term--a comparative study. 2003 *South African Medical Journal.* pp. 371-374.
28. Rehan-Uddin K, El-Refaey H, Sharma S, et al. Oral, rectal, and vaginal pharmacokinetics of misoprostol. *Obstet Gynecol* 2004; 103:866-70.
29. Rouzi AA, Alsibiani S, Mansouri N, Alsanani N, Darhouse K. Randomized clinical trial between hourly titrated oral misoprostol and vaginal dinoprostol for induction of labor. *Am J Obstet Gynecol* 2014; 210:56.e1-6.
30. Rouzi AA, Alsahly N, Alamoudi R, Almansouri N, Alsinani N, Alkafy S, Rozzah R, Abduljabbar H. Randomized clinical trial between hourly titrated and 2 hourly static oral misoprostol solution for induction of labor. *Am J Obstet Gynecol.* 2016 Dec 14
31. Souza AS, Feitosa FE, Costa AA, Pereira AP, Carvalho AS, Paixão RM, Katz L, Amorim MM. Titrated oral misoprostol solution versus vaginal misoprostol for labor induction. *Int J Gynaecol Obstet.* 2013 Dec;123(3):207-12.
32. Souza AS, Scavuzzi A, Rodrigues DC, Oliveira RD, Feitosa FE, Amorim MM. Titrated oral solution of misoprostol for labour induction: a pilot study. *Rev Bras Ginecol Obstet.* 2010 May;32(5):208-13. [Article in Portuguese]
33. Sciscione AC, Nguyen L, Manley J, Pollock M, Maas B, Colmorgen G. A randomized control trial of transcervical Foley catheter to intravaginal misoprostol for preinduction cervical ripening. *Obstet Gynecol* 2001; 97:603-7.
34. Tang OS, Schweer H, Sebberth HW, et al. Pharmacokinetics of different routes of administration of misoprostol. *Hum Reprod* 2002; 17:332-6.
35. Thaisomboon A, Russameecharoen K, Wanitpongpan P, Phattanachindakun B, Changnoi A. Comparison of the efficacy and safety of titrated oral misoprostol and a conventional oral regimen for cervical ripening and labor induction. *Int J Gynaecol Obstet.* 2012 Jan;116(1):13-6. doi: 10.1016/j.ijgo.2011.07.027. Epub 2011 Sep 28.

- Vogel JP, West HM, Dowswell T. Titrated oral misoprostol for augmenting labour to improve maternal and neonatal outcomes. *Cochrane Database Syst Rev*. 2013 Sep 23;(9):CD010648. doi: 10.1002/14651858.CD010648.pub2. (Accessed 4/15/23).
36. Wang X, Yang A, Ma Q, Li X, Qin L, He T. Comparative study of titrated oral misoprostol solution and vaginal dinoprostone for labor induction at term pregnancy. *Arch Gynecol Obstet*. 2016 Jan 8.
 37. Weeks A, Alfirevic Z, Faundes A, Hofmeyr GJ, Safar P, Wing D. Misoprostol for induction of labour with a live fetus. *Int J Gynecol Obstet* 2007; 99:S194-7.
 38. Zvandasara P, Saungweme G, Mlambo J, Chidembo W, Madzivanzira N, et al. Induction of labour with titrated oral misoprostol suspension. A comparative study with vaginal misoprostol. *Central African Journal of Medicine*. (2008) pp. 43-49.
 39. Zeiman M, Fong SK, Benowitz NL, et al. Absorption kinetics of misoprostol with oral or vaginal administration. *Obstet Gynecol* 1997; 90:88-92.

VERSION HISTORY

Revised 5/17/26 pca
Revised 4/15/23 njm
Revised 4/16/21 njm
Revised 5/31/17 njm
Written 3/28/17njm

TABLES**TABLE 1: BISHOP SCORE**

Cervical Exam	Points			
	0	1	2	3
Dilation (cm)	Closed	1-2cm	3-4cm	5-6cm
Effacement (%)	0-30%	40-50%	60-70%	80%
Station	-3	-2	-1, 0	+1, +2
Consistency	Firm	Medium	Soft	
Position	Posterior	Mid	Anterior	

TABLE 2: CERVICAL RIPENING – ORAL TITRATED MISOPROSTOL PROTOCOL

Dose	Frequency	Number of Doses	Fetal Monitoring
20 mcg in 20 mL	Every one hour	Four	20 minutes pre-dose 20 minutes post-dose
40 mcg in 40 mL	Every two hours	Four	20 minutes pre-dose 20 minutes post-dose
60 mcg in 60 mL	Every two hours	Four	20 minutes pre-dose 20 minutes post-dose

TABLE 3: AUGMENTATION – ORAL TITRATED MISOPROSTOL PROTOCOL

Dose	Frequency	Number of Doses	Fetal Monitoring
5 mcg in 5 mL	Every one hour	One	20 minutes pre-dose 20 minutes post-dose
10 mcg in 10 mL	Every one hour	One	20 minutes pre-dose 20 minutes post-dose
20 mcg in 20 mL	Every one hour	Until active labor	20 minutes pre-dose 20 minutes post-dose

FIGURES

FIGURE 1: LEVEL OF PLASMA MISOPROSTOL CONCENTRATION OVER TIME (ZEIMAN, ET AL).

