

Fever in Well-Appearing Infants 3-60 Days Old

Scope and Definitions:

This guideline applies to previously healthy, term (≥ 37 weeks), well-appearing infants aged 3–60 days presenting with fever $\geq 38.0^{\circ}\text{C}$ (100.4°F), without an obvious source of infection. It does not apply to ill-appearing infants, preterm infants, or those with significant comorbidities, who require immediate full sepsis evaluation and empiric treatment regardless of age.

For infants < 3 days old, please refer to NICU Early onset sepsis/Late onset sepsis or NEC guidelines.

Epidemiology and Risk:

- More than 2% of all US full-term infants are evaluated for fever in the first months of life.
- Approximately 8–13% of febrile young infants have a bacterial infection, predominantly urinary tract infections (UTIs).
- Approximately 1–4% of febrile infants ≤ 28 days old harbor invasive bacterial infections (bacteremia and/or bacterial meningitis).
- Infants ≤ 21 days old carry the highest risk of serious bacterial infection (SBI).
- Most febrile young infants with invasive bacterial infections are well-appearing and cannot be reliably identified by history and physical examination alone.

Common Pathogens:

- Group B Streptococcus (GBS): Most common Gram-positive pathogen; responsible for $> 20\%$ of bacteremia after the first week of life and the most common cause of bacterial meningitis.
- Escherichia coli: Most common overall pathogen; accounts for 70–90% of UTIs, 30–60% of bacteremia, and 15–30% of bacterial meningitis.
- Other: Listeria monocytogenes (rare), Enterococcus, Staphylococcus aureus, Streptococcus pneumoniae, Klebsiella species.
- Herpes simplex virus (HSV): Population-based rates of 2–5 per 100,000 neonates; 15% present with fever as the only symptom.

Exclusion Criteria:

- < 37 weeks gestation at birth
- Infants < 2 weeks old with perinatal course complicated by fever/infection/antimicrobial use
- High suspicion for HSV infection (see specific HSV page 5)
- Focal bacterial findings
- Documented or suspected immune compromise
- Complicated neonatal course (surgery or infection)
- Congenital anomalies
- Medically fragile/technology dependent infants

Fever After Immunizations:

Fever is a common and expected adverse event following routine immunizations. In infants younger than 60 days of age, fever occurring within 24 to 48 hours after vaccination may be vaccine-related; however, a serious bacterial infection cannot be excluded based on recent immunization alone. Clinical evaluation should consider the infant's age, appearance, timing of vaccination, temperature, and other risk factors. Recent immunization does not eliminate the need for medical assessment of a febrile infant when clinically indicated.

Table of Contents:

ANMC Fever in Well Appearing Infant 3-21 Days Guideline.....	2
ANMC Fever in Well Appearing Infant 22-28 Days Guideline.....	3
ANMC Fever in Well Appearing Infant 29-60 Days Guideline.....	4
ANMC Fever in Well Appearing Infant HSV	5

ANMC Fever in Well Appearing Infant 3-21 Days Guideline

Temperature $\geq 100.4^{\circ}\text{F}$ (38°C)
from any source, including home thermometer
OR
Temperature $<96.8^{\circ}\text{F}$ (36°C) that does not quickly resolve
with environmental measures

Laboratory Work Up

Blood	Urine	CSF*	HSV Eval – if increased risk ^{††}	Other
• Blood Culture • CBC • CRP • ALT • AST	• Urine culture (cath specimen) • Urinalysis	• CSF culture • Gram stain • Cell count • Glucose • Protein • Multiplex PCR	• Serum HSV PCR • CSF HSV PCR (included in multiplex PCR) • Surface Swab HSV PCR (conjunctivae, nasopharynx, mouth, anus) • CMP	• RSV/Flu/Covid swab if clinically appropriate • Consider stool studies if diarrhea • Consider CXR if indicated

*If unable to obtain LP: re-attempt in 12-24h, treat with meningitic dosing/antibiotics while awaiting re-attempt.

^{††} Maternal history of genital HSV lesions or fevers from 48 hours before to 48 hours after delivery. Infants with vesicles, seizures, hypothermia, mucous membrane ulcers, CSF pleocytosis in the absence of a positive Gram stain result, leukopenia, thrombocytopenia, or elevated alanine aminotransferase levels.

Laboratory Interpretation

Urine: Negative UA does not rule out UTI. Diagnose UTI with urine culture $>10,000$ cfu/mL of single uropathogen

CSF: Normal CSF WBC <20 , Absence of neutrophils (polys) makes bacterial meningitis unlikely; CSF neutrophils (polys) $>75\%$ increases likelihood of bacterial meningitis. If LP difficult to interpret/bloody: Do not use correction formulas for WBC.

No

Suspicion for meningitis?

Yes

Empiric Antibiotics – No Suspicion for Meningitis

Goal: administer within 60 minutes of presentation

Ampicillin

50mg/kg/dose IV q8hrs

AND

Gentamicin

≤ 7 days 4mg/kg/dose IV q24hrs

8-28 days 5mg/kg/dose IV q24hrs

ADD IF APPROPRIATE

Acyclovir

20mg/kg/dose IV q8hrs

Empiric Antibiotics – Suspicion for Meningitis

Goal: administer within 60 minutes of presentation

Consider ID Consultation

Ampicillin – Meningitis Dosing

≤ 7 days 100mg/kg/dose IV q8hrs

8-28 days 75mg/kg/dose IV q6hrs

AND

Cefepime

50mg/kg/dose IV q8hrs

ADD IF APPROPRIATE

Acyclovir

20mg/kg/dose IV q8hrs

Admit to the hospital

Pathogen or source identified at 24-36 hours?

No

If all cultures and PCRs negative. Stop antimicrobials.
Discharge if clinically appropriate.

Yes

If source identified, treat infection.
If only UTI, consider transition to oral therapy.

ANMC Fever in Well Appearing Infant 22-28 Days Guideline

Temperature $\geq 100.4^{\circ}\text{F}$ (38°C) from any source, including home thermometer
OR
 Temperature $\leq 96.8^{\circ}\text{F}$ (36°C) that does not quickly resolve with environmental measures

Laboratory Work Up

Blood	Urine	Other
-Blood Culture -CBC -CRP	-Urine culture (cath specimen) -Urinalysis	-RSV/Flu/Covid swab if clinically appropriate -consider stool studies if diarrhea -consider CXR if indicated

- Urine: Negative UA does not rule out UTI. Diagnose UTI with urine culture $>10,000\text{cfu/mL}$ of single uropathogen.
- Abnormal inflammatory marker: CRP $>2\text{ mg/dL}$, ANC $>4000\text{ neutrophils}/\mu\text{L}$

Abnormal UA OR
 ≥ 1 abnormal
 inflammatory marker

Perform LP (CSF culture,
 cell count, PCR panel,
 Protein, Glucose)

Normal UA AND
 Normal inflammatory
 markers

No LP,
 No antibiotics

Laboratory Interpretation

CSF: Normal CSF WBC <15 , Absence of neutrophils (polys) makes bacterial meningitis unlikely; CSF neutrophils (polys) $>75\%$ increases likelihood of bacterial meningitis. If LP difficult to interpret/bloody: Do not use correction formulas for WBC.

No

Suspicion for meningitis?

Yes

Empiric Antibiotics – No Suspicion for Meningitis

Goal: administer within 60 minutes of presentation

Ampicillin

50mg/kg/dose IV q8hrs

AND

Gentamicin

5mg/kg/dose IV q24hrs

Empiric Antibiotics – Suspicion for Meningitis

Goal: administer within 60 minutes of presentation.

Consider ID consultation.

Ampicillin – Meningitis Dosing

75mg/kg/dose IV q6hrs

AND

Cefepime

50mg/kg/dose IV q8hrs

Pathogen or source identified at 24-36 hours?

No

Yes

If all cultures and PCRs negative. Stop antimicrobials.
 Discharge if clinically appropriate.

If source identified, treat infection.
 If only UTI, consider transition to oral therapy.

ANMC Fever in Well Appearing Infant 29-60 Days Guideline

Temperature $\geq 100.4^{\circ}\text{F}$ (38°C) from any source, including home thermometer
OR
Temperature $\leq 96.8^{\circ}\text{F}$ (36°C) that does not quickly resolve with environmental measures

Clinical suspicion of
bronchiolitis?

Yes

Laboratory Work Up

Urine	Other
-Urine culture (cath specimen)	-RSV/Flu/Covid PCR (+/- Pertussis if circulating)
-Urinalysis	-consider blood culture and CBC
	-consider CXR if indicated

No

Laboratory Work Up

Blood	Urine	Other
-Blood Culture	-Urine culture (cath specimen)	-consider stool studies if diarrhea
-CBC	-Urinalysis	-consider CXR if indicated
-CRP		

- Urine: Negative UA does not rule out UTI. Diagnose UTI with urine culture $>10,000\text{cfu/mL}$ of single uropathogen.
- Abnormal inflammatory marker: CRP $>2\text{ mg/dL}$, ANC $>4000\text{ neutrophils}/\mu\text{L}$

Laboratory Interpretation

CSE: Normal CSF WBC <9 , Absence of neutrophils (polys) makes bacterial meningitis unlikely; CSF neutrophils (polys) $>75\%$ increases likelihood of bacterial meningitis. If LP difficult to interpret/bloody: Do not use correction formulas for WBC.

Normal UA,
Abnormal
inflammatory
markers

Abnormal UA,
Abnormal
inflammatory
markers

Abnormal UA,
Normal
inflammatory
markers

Normal UA AND
Normal
inflammatory
markers

Perform LP (CSF
culture, cell count, PCR
panel, Protein, Glucose)

LP is a "weak
recommendation" if any
inflammatory marker is
abnormal

No LP

Cephalexin
17mg/kg/dose PO TID
x7-10 days

No LP,
No antibiotics

No

Suspicion for meningitis?

Yes

Empiric Antibiotics – Suspicion for Meningitis

Goal: administer within 60 minutes of presentation.
Consider ID Consultation.

Ceftriaxone – Meningitis Dosing

50mg/kg/dose IV q12hrs

AND

Vancomycin

15mg/kg/dose IV q6-8hrs (depending on post-menstrual age), Pharmacy to dose

Empiric Antibiotics – No Suspicion for Meningitis

Goal: administer within 60 minutes of presentation

Ceftriaxone

50mg/kg/dose IV q24hrs

Pathogen or source identified at 24-36 hours?

No

Yes

If all cultures and PCRs negative. Stop antimicrobials.
Discharge if clinically appropriate.

If source identified, treat infection.
If only UTI, consider transition to oral therapy.

ANMC Fever in Well Appearing Infant HSV

Age	HSV Risk Factors	HSV Studies	HSV Treatment
<6 weeks old	• Toxic or critically ill appearance (not included in this guideline) • Prematurity: <37 weeks EGA - off guideline • Known or suspected HSV exposure • Vesicles or mucous membrane ulcers • Seizure • Persistent hypothermia: <36°C • Thrombocytopenia: <150/mm³ • Elevated alanine aminotransferase (if obtained) >1.5x ULN	• HSV blood PCR • HSV CSF PCR • HSV cutaneous lesions PCR – unroof one or more vesicular lesions and obtain vesicular fluid on an appropriate swab • HSV surface swabs PCR – swab conjunctivae, nasopharynx, mouth, anus; may be obtained as a single swab starting with the conjunctivae and moving caudally • Liver function tests (AST, ALT)	Acyclovir 20mg/kg/dose IV q8h
>6 weeks old	• Vesicles or mucous membrane ulcers • Risk of Neonatal HSV is highest in the first four weeks, and is rare at and beyond 6 weeks of life.	• HSV cutaneous lesions PCR – unroof one or more vesicular lesions and obtain vesicular fluid on an appropriate swab	

References

1. Pantell RH, Roberts KB, Adams WG, et al; Subcommittee on Febrile Infants. Evaluation and management of well-appearing febrile infants 8 to 60 days old. *Pediatrics*. 2021;148(2):e2021052228. doi:10.1542/peds.2021-052228. [\[publications.aap.org\]](https://publications.aap.org)
2. American Academy of Pediatrics. Infant fever: clinical practice guideline for evaluation and management of well-appearing febrile infants 8-60 days old. Accessed September 4, 2026. <https://www.aap.org/en/patient-care/infant-fever/> [\[aap.org\]](https://www.aap.org/en/patient-care/infant-fever/)
3. Kuppermann N, Dayan PS, Levine DA, et al. A Clinical Prediction Rule to Identify Febrile Infants 60 Days and Younger at Low Risk for Serious Bacterial Infections. *JAMA Pediatr*. 2019;173(4):342-351. doi:10.1001/jamapediatrics.2018.5501
4. Northern California Pediatric Hospital Medicine Consortium. Consensus guidelines for febrile infants 0-90 days of age. University of California San Francisco Benioff Children's Hospitals. Updated August 10, 2023. Accessed September 4, 2026. <https://medconnection.ucsfbenioffchildrens.org/febrile-infant-guidelines> [\[medconnect...ldrens.org\]](https://medconnect...ldrens.org)
5. Thompson S, Kenneally A, et al. Fever in well-appearing infants less than 60 days: clinical pathway. Riley Children's Health; updated February 2025. Accessed September 4, 2026. <https://cdn.rileychildrens.org/resources/Clinical-Pathways-and-Guidelines/Fever-in-Well-Appearing-Infants-Less-than-60-Days-Old.pdf>
6. Mahajan P, Browne LR, Levine DA, Cohen DM, Gattu R, Linakis JG, Anders J, Borgialli D, Vitale M, Dayan PS, Casper TC, Ramilo O, Kuppermann N; Febrile Infant Working Group of the Pediatric Emergency Care Applied Research Network (PECARN). Risk of Bacterial Coinfections in Febrile Infants 60 Days Old and Younger with Documented Viral Infections. *J Pediatr*. 2018 Dec;203:86-91.e2. doi: 10.1016/j.jpeds.2018.07.073. Epub 2018 Sep 6. PMID: 30195552; PMCID: PMC7094460.
7. Melvin A, Mohan K. Neonatal Herpes Simplex Virus Infection: Epidemiology and Outcomes in the Modern Era, *Journal of the Pediatric Infectious Diseases Society*, 2021;:, piab105, <https://doi.org/10.1093/jpids/piab105>